

A History of Abuse and Operative Delivery – Results from a European Multi-Country Cohort Study

Berit Schei^{1,2}, Mirjam Lukasse^{1,3*}, Elsa Lena Ryding⁴, Jacquelyn Campbell⁵, Helle Karro⁶, Hildur Kristjansdottir^{7,8}, Made Laanpere⁶, Anne-Mette Schroll⁹, Ann Tabor^{9,10}, Marleen Temmerman¹¹, An-Sofie Van Parys¹¹, Anne-Marie Wangel¹², Thora Steingrimsdottir^{7,13}

1 Department of Public Health and General Practice, Norwegian University of Science and Technology, Trondheim, Norway, **2** Department of Obstetrics and Gynaecology, St.Olav's University Hospital, Trondheim, Norway, **3** Department of Health, Nutrition and Management, Oslo and Akershus University College of Applied Sciences, Oslo, Norway, **4** Department of Women's and Children's Health, Division of Obstetrics and Gynaecology, Karolinska Institutet/University Hospital, Stockholm, Sweden, **5** John Hopkins University, School of Nursing, Baltimore, Maryland, United States of America, **6** Department of Obstetrics and Gynaecology, University of Tartu, Tartu, Estonia, **7** Department of Obstetrics and Gynaecology, Landspítali University Hospital, Reykjavik, Iceland, **8** Directorate of Health, Reykjavik, Iceland, **9** Centre of Fetal Medicine, Department of Obstetrics, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark, **10** Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark, **11** Department of Obstetrics and Gynaecology, Ghent University Hospital, Ghent, Belgium, **12** Malmö University, Faculty of Health and Society, Malmö, Sweden, **13** Primary Health Care of the Capital Area, Centre of Development, Reykjavik, Iceland

Abstract

Objective: The main aim of this study was to assess whether a history of abuse, reported during pregnancy, was associated with an operative delivery. Secondly, we assessed if the association varied according to the type of abuse and if the reported abuse had been experienced as a child or an adult.

Design: The Bidens study, a cohort study in six European countries (Belgium, Iceland, Denmark, Estonia, Norway, and Sweden) recruited 6724 pregnant women attending routine antenatal care. History of abuse was assessed through questionnaire and linked to obstetric information from hospital records. The main outcome measure was operative delivery as a dichotomous variable, and categorized as an elective caesarean section (CS), or an operative vaginal birth, or an emergency CS. Non-obstetrically indicated were CSs performed on request or for psychological reasons without another medical reason. Binary and multinomial regression analysis were used to assess the associations.

Results: Among 3308 primiparous women, sexual abuse as an adult (≥ 18 years) increased the risk of an elective CS, Adjusted Odds Ratio 2.12 (1.28–3.49), and the likelihood for a non-obstetrically indicated CS, OR 3.74 (1.24–11.24). Women expressing current suffering from the reported adult sexual abuse had the highest risk for an elective CS, AOR 4.07 (1.46–11.3). Neither physical abuse (in adulthood or childhood < 18 years), nor sexual abuse in childhood increased the risk of any operative delivery among primiparous women. Among 3416 multiparous women, neither sexual, nor emotional abuse was significantly associated with any kind of operative delivery, while physical abuse had an increased AOR for emergency CS of 1.51 (1.05–2.19).

Conclusion: Sexual abuse as an adult increases the risk of an elective CS among women with no prior birth experience, in particular for non-obstetrical reasons. Among multiparous women, a history of physical abuse increases the risk of an emergency CS.

Citation: Schei B, Lukasse M, Ryding EL, Campbell J, Karro H, et al. (2014) A History of Abuse and Operative Delivery – Results from a European Multi-Country Cohort Study. PLoS ONE 9(1): e87579. doi:10.1371/journal.pone.0087579

Editor: Shannon M. Hawkins, Baylor College of Medicine, United States of America

Received: September 24, 2013; **Accepted:** December 23, 2013; **Published:** January 31, 2014

Copyright: © 2014 Schei et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The Bidens study was supported by the Daphne II Program, European Commission for Freedom, Security, and Justice, Brussels, Belgium (Grant no. JLS/2006/DAP-1/242/W30-CE-0120887/00-87). Mirjam Lukasse received a postdoctoral fellowship from the Norwegian Research Council, Grant no. 204292. The funders had no role in study design, data collection and analysis, decision to publish, or presentation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Mirjam.Lukasse@Hioa.no

Introduction

Interventions during childbirth, such as operative delivery, aim to ensure best possible health for the mother and child. There are interventions that are obviously necessary and life-saving for both mother and child [1]. However, sometimes the indication for intervention is subject to debate and even considered non-medical, such as caesarean section (CS) performed for fear of childbirth or on maternal request only [2,3].

Violence against women (VAW) is a broad term covering a large range of traumatic events and abuse occurring during a woman's life-span. VAW has been shown to increase the risk for complications during pregnancy and may thereby increase interventions during delivery [4–8]. VAW also increases the risk of psychological distress and fear of childbirth, which in turn may become the indication for CS on maternal request [9–11]. Furthermore, VAW may alter a woman's stress response and affect ability to communicate with obstetric staff during labor, thus

increasing the risk of interventions [12]. Few studies exist which examine the association between abuse and mode of delivery. The studies that exist are limited in terms of the type of abuse addressed and inconclusive in their findings [13–18]. Two studies show a major increase in risk of CS for women who have been subjected to rape in adulthood [15,18]. However, the women included were referred to a mental health team for known psychological problems and because of their desire to give birth by elective CS [15,18]. The association between abuse and mode of delivery may not be the same among women attending routine antenatal care. The aim of this study was to test the a priori hypothesis that a history of abuse is associated with an increased risk of operative delivery, i.e. a caesarean section (elective or emergency) or an operative vaginal delivery, among women attending routine antenatal care. Secondly, we wanted to assess if potential associations varied between types of abuse (emotional, physical or sexual), and whether the abuse had occurred during childhood or adulthood. Thirdly, where an association between a type of abuse and CS was found, we explored the association with non-obstetric indication.

Methods

The Bidens study, a six-country (Belgium, Iceland, Denmark, Estonia, Norway, and Sweden) cohort study of unselected pregnant women, was the result of an EU-funded collaboration between the Norwegian University of Science and Technology (NTNU) and partners from Universities and Hospitals in six European countries. A short description of the study sites is given in Table 1.

Participants

Recruitment took place between March 2008 and August 2010. Women who consented subsequently completed a questionnaire and allowed the extraction of specified data on their delivery from their medical notes. Due to country specific organization as well as requirements of local ethical committees, minor variations in the recruitment procedure occurred.

In *Belgium*, women were approached by the midwife or secretary when attending antenatal care. Consenting women were asked to complete the questionnaire in a separate room. In *Iceland* women were recruited when attending routine ultrasound and returned completed forms by mail. In *Denmark*, women were given information about the study when attending early routine ultrasound screening and were mailed the questionnaire later. They returned the questionnaire by mail or when attending their next ultrasound examination. In *Estonia*, women were invited to participate while visiting for an antenatal consultation. After completing the questionnaire, it was left in a mailbox at the clinic. In *Norway* women, after attending routine ultrasound, received the questionnaire by mail and returned it by mail. Non-responders were sent one reminder. In *Sweden*, the questionnaire was administered to women when attending routine glucose tolerance test and filled out during the two hours gap between the blood samplings. Belgium and Sweden were not permitted to record non-participation. The estimated response rate varied between 50% in Norway to 90% in Estonia.

Inclusion and exclusion criteria

All women required sufficient language skills to fill out the form. In Estonia, women could choose to fill out an Estonian or Russian

Table 1. Presentation of recruitment sites in the Bidens study.

Country	City	Characteristics (inhabitants)	Recruitment Site	Number of deliveries in 2008	Number of deliveries in geographical area
Belgium	Gent	Third largest city (240.000)	Gent University Hospital	1217	6674
	Waregem	(36.000)	Onze Lieve Vrouw van Lourdes/'Our Lady from Lourdes'	721	721
	leper	(35.000)	Jan Yperman Hospital	1091	1091
Iceland	Reykjavik	Capital (200.000)	Landspítali University Hospital	3373	4118
Denmark	Copenhagen	Capital (1.811.239)	University Hospital	3730	21101§
Estonia	Tallinn	Capital (400 000)	West-Tallinn Central Hospital	3283	7421
			East-Tallinn Central Hospital	4386	
	Tartu	Second largest city in Estonia (100 000)	Tartu University Hospital	2325	1994
	Kohtla-Järve	North-East Estonia (46 000) 80% Russian speaking	East-Viru Central Hospital	515	1490
Norway	Drammen	(60 000)	Buskerud Regional Hospital	1961	3003*
	Oslo	Capital (560 000)	Rikshospitalet, OUS	2238	10252*
	Tromsø	Most northern city (67 000)	University hospital in North -Norway	1509	1961*
	Trondheim	(165 000)	St.Olavs University hospital	3483	3830*
	Ålesund	(42 000)	Hospital in Ålesund	1291	2813*
Sweden	Malmö	(295 000)	Antenatal Care Clinics (ANC):	University Hospital MAS 4359	Selected ANC represent Approx. 60% of all births of the catchments area

References for number of deliveries in the geographical area: Belgium: SPE (Studiecentrum Perinatale Epidemiologie) 2008. Iceland: The Icelandic Birth register for 2008. Denmark: http://www.sst.dk/Indberetning%20og%20statistik/Sundhedsdata/Foedsler_fertiliteetsbehandling_og_abort/foedsler1.aspx.

§Born at hospital Estonia: <http://www.tai.ee/et/tegevused/registrid/meditsiiniline-sunniregister-ja-raseduskatkestus-andmekogu/statistika>; Estonian Medical Birth Registry.

Sweden regional data: <http://www.socialstyrelsen.se/register/halsodataregister/medicinskafodelseregistret> Norway: Medical Birth Registry.

*Number of newborn ≥ 22 weeks gestation.

doi:10.1371/journal.pone.0087579.t001

language questionnaire. In Belgium, Iceland and Denmark women less than 18 years of age were excluded. In Denmark, only women from the local geographical area were invited. In Belgium, women who could not be separated from their accompanying person were not recruited. In Iceland, Denmark and Norway, women with major fetal pathology were excluded from the study.

Sample size for this study. Of the 7200, women who consented and returned the questionnaire 6724 were included in this study. Of the 476 excluded, 47 had failed to answer two or more of the questions on abuse, 304 lacked information about mode of delivery, 122 lacked parity, and the pregnancy of three women ended before 22 weeks gestation.

Questionnaire

The main instrument of the present study is a 68 items questionnaire, partly based on the NorAQ (Norvold Abuse Questionnaire, Figure 1), which was developed in a previous multi-centre study among gynaecological patients in the Nordic countries [19]. The different types of abuse and severity of abuse were defined in NorAQ by a validated set of thirteen descriptive questions [20]. Also included in the questionnaire were questions on post-traumatic stress symptoms [21], fear of childbirth, using the Wijma Delivery Expectancy Questionnaire (W-DEQ) [22], and a short version of the Edinburgh Postpartum Depression Scale (EPDS-5) [23]. Method and experience of previous deliveries, as well as preference of mode of delivery, were assessed. A complete version of the questionnaire was developed in English. Where a previously translated version of the NorAQ, W-DEQ or EPDS was available, this was used. Otherwise, the questionnaire was translated into the required languages by a native speaker (Flemish, Icelandic, Danish, Estonian, Russian, Norwegian and Swedish) and then translated back again into the source language. The original and back-translated versions were used to determine the final version.

Follow-up

Birth outcome data was collected from hospital records and recorded on the outcome sheet prepared for this study.

Variables

Main exposure: Emotional, physical and sexual abuse was assessed in three identically structured sections [19]. For each type and level of abuse, the answer categories were no, yes as a child, yes as an adult, or yes both as a child and as an adult and classified according to the most severe level reported (mild, moderate and, severe). Women were asked if they experienced the indicated abuse during the past 12 months, which was coded as recent. The degree of current suffering was measured on a visual analogous scale (0–10) and recoded into zero for values 0–4 and 1 for values 5–10. Women were defined as having experienced any abuse if they answered yes to at least one of the questions of sexual, emotional and physical abuse, except mild physical abuse, which showed low specificity in the validation study [20].

Education was coded into four levels: primary school (9 years), secondary school (13 years), higher education (university or college), <4 years and ≥ 4 years. Women were considered to have a twin pregnancy if they had reported this in the questionnaire.

The outcome, operative delivery was defined as a dichotomous yes or no variable to test the main hypothesis for this study, and as a categorical variable with the following categories: 0) spontaneous vaginal 1) elective Caesarean Section (CS) 2) forceps or vacuum extraction and 3) emergency CS. Indications options on the outcome sheet for operative deliveries included: fetal distress,

dystocia, breech, maternal exhaustion, maternal request, psychosocial indications, other obstetrical indications, and unknown. A CS was defined as non-obstetrically indicated when “maternal request” or “psychosocial indication” were the reported reasons without another medical indication. The indication for the operative delivery was taken from the hospital record, as was gestational age (based on ultrasound during pregnancy) at birth.

Ethical considerations

The study was conducted in accordance with the ethical guidelines developed by WHO [24]. The information letter instructed women to complete the form in a place where they could be undisturbed and included telephone numbers and e-mail addresses to contact if needed. Additionally, in Belgium, Estonia and Sweden the participants had the opportunity to complete the questionnaires at the clinic, and measures were installed to avoid accompanying persons to be with them. Formal approvals of local ethical committees and data protection agencies were obtained at all sites, as listed below.

Belgium: The Ethical Committee of Ghent University acted as the central ethical committee for the study; U(Z) Gent, 22012008/B67020072813, date of approval: 1st February 2008, Waregem hospital date added: 21st October 2008.

Iceland: The scientific board approved the study (24.06.2008-VSN-b2008030024/03-15) according to Icelandic regulations, date: 24th June 2008.

In *Denmark*, even though ethical approval for non-invasive studies is not required, the study was presented to the Research Ethics Committee of the Capital Region, who found no objections to the study (H-A-2008-002), date: 11th February 2008. Permission was obtained from the Danish Data Protection Agency (J.nr. 2007-41-1663).

In *Estonia*, ethical permission was given by the Ethics Review Committee on Human Research of the University of Tartu, Estonia; 190/M-29, 192/-22, 196/X-2, date: 17th December 2007, East-Tallinn Central Hospital added: 19th January 2009, Russian language and prolonged period added: 22nd February 2010, East-Viru Central Hospital added: 26th April 2010.

In *Norway*, the Regional Committee for Medical Research Ethics in North approved the study (72/2006), date: 29th August 2007; and the Data Inspectorate (NSD) (15214/3/) also approved the study, date: 19th December 2007.

In *Sweden*, the study was approved by the Regional Ethical Committee in Stockholm (2006/354-31/1), date: 14th June 2006.

The data was anonymised prior to analysis.

Statistical analyses

Power calculation was based on the main hypotheses that exposure to any abuse increased the risk of any operative delivery, OR 1.4 ($\alpha = 0.05$, $\beta = 0.20$), assuming one exposed and four non-exposed within an unselected cohort of pregnant women [19,25]. In total, 2500 women needed to be recruited, allowing for stratified analysis based on parity approximately 5000.

Cross-tabulation was used to quantify socio-demographic, life-style and obstetric characteristics factors by country of residence and mode of delivery. Frequency analyses were used to quantify the prevalence of the different types of abuse by category, age <18 or ≥ 18 , level of severity, current suffering and recentness. All regression analyses were stratified for parity. The main hypothesis was tested by binary logistic regression analysis. The association between different abuse categories and the different kinds of operative delivery were analysed by multinomial regression analyses. Based on the literature and our experience, we included all of the variables (which we had information on) correlated with

Level of abuse	Type of abuse
	Emotional
Mild	Have you experienced anybody systematically and for a long period trying to repress, degrade or humiliate you?
Moderate	Have you experienced anybody and by threat or force trying to restrict your contacts with others or totally control what you may or may not do?
Severe	Have you experienced living in fear because somebody systematically and for a long period has threatened you or somebody close to you?
	Physical
Mild	Have you experienced anybody hitting you, smacking your face, or holding you firmly against your will?
Moderate	Have you experienced anybody hitting you with his/her fist(s) or with a hard object, kicking you, pushing you violently, giving you a beating, thrashing you, or doing anything similar to you?
Severe	Have you experienced anybody threatening your life by, for instance, trying to strangle you, showing a weapon or knife, or by any other similar act?
	Sexual
Mild, no genital contact	Has anybody against your will touched parts of your body other than the genitals in a sexual way or forced you to touch other parts of his or her body in a sexual way?
Mild, emotional or sexual humiliation	Have you in any other way been humiliated; eg. by being forced to watch a pornographic film or similar against your will, forced to participate in a pornographic film or similar, forced to show your body naked, or forced to watch when somebody else showed his/her body naked?
Moderate, genital contact	Has anybody against your will touched your genitals, used your body to satisfy him/herself sexually, or forced you to touch anybody else's genitals?
Severe, penetration	Has anybody against your will put or tried to put his penis into your vagina, mouth, or rectum; put or tried to put an object or other part of the body into your vagina, mouth or rectum?
Answer categories	1=no; 2=yes, as a child (<18years); 3=yes, as an adult (≥18 years); 4=yes, as a child and as an adult

Figure 1. The Norvold Abuse Questionnaire (NorAQ) questions on emotional, physical and sexual abuse.
doi:10.1371/journal.pone.0087579.g001

operative delivery (outcome) and, for each of them evaluated if they were likely to be the result of the exposure (abuse). Of the factors influencing operative delivery we considered the following as correlated to but not the result of the exposure: age, twin pregnancy, gestational age at birth, and country of residence. These variables were included in the adjusted regression analyses. Smoking status [25,26], alcohol consumption [26,27], use of epidural analgesia [14,18], birth weight [26,27] could be the result of the independent variable (abuse) and were not included in the model. Factors such as depressive symptoms, post-traumatic stress symptoms and fear of childbirth were also excluded from the model as we considered them to fall on the pathway between exposure (abuse) and operative delivery. As a previous CS could be the result of abuse before the related pregnancy, we did not want to enter it into all the analyses, but estimated its impact on the significant association(s) found for multiparous women. We estimated crude and adjusted odd ratios (AORs) and 95% confidence intervals (CIs). All analyses were two-sided at $\alpha=0.05$. The comparison group for all analyses was women without a reported history of abuse. The statistical program used was PASW 20.

Results

Among the 6724 women in our study, 2323 (34.5%) reported having experienced any abuse (of any type) at some point in their lives, 1567 (23.3%) as a child, 1309 (19.5%) as an adult and 553 (8.2%) both as a child and an adult. The distribution of responses between countries, for socio-demographic and obstetrical characteristics are presented in Table 2. The participants in Denmark (both primiparous and multiparous women) were older than women elsewhere, while the youngest primiparous women were in Iceland and multiparous in Belgium. The caesarean section (CS) rate was highest in Denmark and lowest in Sweden. Birth weight below 2500 g was highest in Denmark and birth weight ≥ 4000 g was most common in Iceland and least common in Belgium. Socio-demographic and obstetric characteristics by mode of delivery can be found in table 3.

A history of any abuse was not associated with any operative delivery (as a dichotomous variable), neither among primiparous or multiparous women delivery, AOR 1.16 (0.99–1.36) and AOR 1.04 (0.86–1.25) respectively (not in the tables). Among multiparous women, only a history of physical abuse was associated with a significant increase in emergency CSs (Table 4), AOR 1.51 (1.05–2.19). This association was attenuated when we added previous CS as a covariate into the analysis, OR 1.48 (1.001–2.18). Of the 512

Table 2. Socio-demographic and obstetric characteristics of women in the Bidens study N = 6724.

	Belgium n = 819	Iceland n = 585	Denmark n = 1268	Estonia n = 874	Norway n = 2234	Sweden n = 944	Total N = 6724
Age mean (SD)	28.4 (4.3)	29.7 (5.0)	31.9 (4.0)	28.2 (5.1)	30.5 (5.0)	30.4 (4.6)	30.1 (4.9)
Primiparous mean (SD)	27.2 (4.0)	26.8 (4.4)	30.7 (4.0)	25.5 (4.2)	28.6 (4.9)	29.1 (4.4)	28.4 (4.7)
Multiparous mean (SD)	29.8 (4.2)	31.6 (4.5)	33.4 (3.6)	30.5 (4.7)	32.0 (4.4)	32.1 (4.4)	31.7 (4.4)
n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Education							
Primary (6–9 years)	13 (1.6)	42 (7.2)	19 (1.5)	69 (7.9)	53 (2.4)	31 (3.3)	227 (3.4)
Secondary (10–13 years)	301 (36.8)	141 (24.1)	118 (9.3)	286 (32.7)	568 (25.4)	278 (29.4)	1692 (25.2)
College/University <4 year	338 (41.39)	184 (31.5)	460 (36.3)	242 (27.7)	688 (30.8)	221 (23.4)	2133 (31.7)
College/University ≥4 years	161 (19.7)	215 (36.8)	660 (52.1)	270 (30.9)	913 (40.9)	401 (42.5)	2620 (39.0)
Missing	6 (0.7)	3 (0.5)	11 (0.9)	7 (0.8)	12 (0.5)	13 (1.4)	52 (0.8)
Civil status							
Married/Cohabiting	784 (95.7)	542 (92.6)	1206 (95.6)	828 (94.7)	2144 (96.0)	904 (95.8)	6408 (95.3)
Other	25 (3.19)	39 (6.7)	49 (3.9)	45 (5.1)	77 (3.4)	37 (3.9)	272 (4.0)
Missing	10 (1.29)	4 (0.7)	13 (1.0)	1 (0.1)	13 (0.6)	3 (0.3)	44 (0.7)
Parity							
Multiparous	380 (46.4)	356 (60.9)	535 (42.2)	469 (53.5)	1276 (57.1)	401 (42.5)	3416 (50.8)
Primiparous	439 (53.6)	229 (39.1)	733 (57.8)	406 (46.5)	958 (42.9)	543 (57.5)	3308 (49.2)
Gestational age at birth							
<37 weeks	42 (5.1)	24 (4.1)	73 (5.8)	46 (5.3)	121 (5.4)	55 (5.8)	361 (5.4)
≥37 weeks	776 (94.7)	561 (95.6)	1194 (94.2)	826 (94.5)	2102 (94.1)	888 (94.1)	6347 (94.4)
Missing	1 (0.1)	0	1 (0.1)	2 (0.2)	11 (0.5)	1 (0.1)	16 (0.2)
Induced	192 (23.4)	117 (20.0)	245 (19.3)	112 (12.8)	408 (18.3)	83 (8.8)	1157 (17.2)
Epidural Analgesia	552 (67.4)	197 (33.7)	387 (30.5)	112 (12.8)	585 (26.2)	181 (19.2)	2014 (30.0)
Mode of delivery							
Spontaneous vaginal	590 (72.0)	439 (75.0)	895 (70.6)	679 (77.7)	1696 (75.9)	735 (77.9)	5034 (74.9)
Operative delivery	229 (28.0)	146 (25.0)	373 (29.4)	195 (22.3)	538 (24.1)	209 (22.1)	1690 (25.1)
Elective CS	71 (8.7)	40 (6.8)	138 (10.9)	63 (7.2)	171 (7.7)	59 (6.3)	542 (8.1)
Operative vaginal	98 (12.0)	57 (9.7)	117 (9.2)	50 (5.7)	189 (8.5)	69 (7.3)	580 (8.6)
Emergency CS	60 (7.3)	49 (8.4)	118 (9.3)	82 (9.4)	178 (8.0)	81 (8.6)	568 (8.4)
Birth weight							
<2500 g	25 (3.1)	17 (2.9)	61 (4.8)	30 (3.4)	80 (3.6)	30 (3.2)	243 (3.6)
2500–4000 g	733 (89.5)	410 (70.1)	1003 (79.1)	680 (77.8)	1756 (78.8)	728 (77.1)	5310 (79.0)
≥4000 g	59 (7.2)	158 (27.0)	196 (15.5)	162 (18.5)	391 (17.5)	183 (19.4)	1149 (17.1)
Missing	2 (80.2)	0	8 (0.6)	2 (0.2)	7 (0.3)	3 (0.3)	22 (0.3)
Apgar score							

Table 2. Cont.

	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
<7 after 5 min.	7 (0.9)	18 (3.1)	4 (0.3)	9 (1.0)	19 (0.9)	17 (1.8)	74 (1.1)		
≥7 after 5 min.	804 (98.2)	562 (96.1)	1253 (98.8)	860 (98.4)	2199 (98.4)	910 (96.4)	6588 (98.0)		
Missing	8 (1.0)	5 (0.9)	11 (0.9)	5 (0.6)	16 (0.7)	17 (1.8)	62 (0.9)		
Twins	19 (2.3)	5 (0.9)	45 (3.5)	14 (1.6)	53 (2.4)	20 (2.1)	156 (2.3)		

doi:10.1371/journal.pone.0087579.t002

(15.0%) multiparous women who had a CS, 76 (14%) had a non-obstetrically indicated CS. No significant association between a history of physical abuse and non-obstetrically indicated CS was observed, crude OR 0.94 (0.52–1.70). The most common reason for emergency CS among multiparous women with a history of physical abuse was fetal distress, followed by other medical reasons and dystocia. Table 5 presents different types of childhood and adult abuse by operative delivery among primiparous women. Women with no prior birth experience who reported a history of adult sexual abuse were significantly more likely to be delivered by elective CS, AOR 2.12 (1.28–3.49). This likelihood increased when they in addition reported either physical or emotional abuse experienced as an adult (Table 6). Primiparous women expressing current suffering from the reported adult sexual abuse had the highest risk for an elective CS, AOR 4.07 (1.46–11.3). Of all the primiparous women, 18% (598) were delivered by CS. Of these, 42 (7.0%) had a non-medical indication. Adult sexual abuse increased the odds of a CS without a medical indication, OR 3.74 (1.24–11.24). The most common indication for an elective CS among primiparous women who reported adult sexual abuse was a breech presentation, followed by other medical reasons and maternal request.

Discussion

In our multi-country study of pregnant women attending routine antenatal care, followed through to delivery, the main hypothesis that a history of any abuse was associated with an operative delivery was not confirmed. Rather, among multiparous women, a reported history of physical abuse was the only type of abuse associated with operative delivery, as it increased the odds for emergency caesarean section. Among primiparous women, only a history of adult sexual abuse was associated with operative delivery, through an increase in elective CS. Primiparous women with a history of adult sexual abuse were more likely to have a non-obstetrical indication when delivered by CS. In addition was current suffering from adult sexual abuse associated with an increased risk for an emergency CS.

A great strength of our study is that it is based on pregnant women attending routine antenatal care. The recruitment procedure varied across sites, both for the invitation to participate, where the questionnaire was completed and how it was returned. This may have introduced information bias. The participation rates varied between countries, but the background characteristics did not indicate any significant selection bias when compared to information from official health authorities. Guidelines for antenatal care and interventions during delivery vary between the participating countries. For example, neither in Belgium nor in Estonia is maternal request alone an accepted indication for medical interventions such as delivery by CS. The uniform results across the participating countries strengthen our finding that among primiparous women a history of sexual abuse as an adult increases the risk for elective CS, regardless of maternal request being an acceptable indication. At all the sites included in the study, epidural analgesia during labour is available, however the rates of use varied greatly between the countries. We do not know how this may influence our results.

Victims of adult sexual abuse with posttraumatic stress symptoms may avoid triggers like participation in our study and have a particular high risk of interventions during delivery. On the other hand, simply answering our questionnaire could be considered an intervention in itself. Recalling abuse when completing the questionnaire may trigger fear of childbirth and symptoms of depression, which again may affect the mode of

Table 3. Socio-demographic and obstetric characteristics of women in the Bidens study (N = 6724) by mode of delivery.

	Elective CS n = 542	Spontaneous vaginal n = 5034	Operative vaginal n = 580	Emergency CS n = 568
Age: mean (SD)	31.9 (5.0)	30.0 (4.8)	29.5 (4.7)	30.1 (5.1)
Primiparous	30.1 (5.3)	28.1 (4.6)	29.1 (4.6)	29.1 (4.9)
Multiparous	32.9 (4.5)	31.6 (4.4)	31.2 (4.6)	32.5 (4.6)
	n (%)	n (%)	n (%)	n (%)
Education				
Primary (6–9 years)	11 (2.1)	182 (3.6)	12 (2.1)	22 (3.9)
Secondary (10–13 years)	135 (25.3)	1275 (25.5)	140 (24.4)	142 (25.2)
College/University <4 year	168 (31.5)	1574 (31.5)	188 (32.8)	203 (36.1)
College/University ≥4 years	219 (41.1)	1972 (39.4)	233 (40.7)	196 (34.8)
Civil status:				
Married/Cohabiting	518 (96.3)	4800 (95.9)	552 (96.3)	538 (95.2)
Other	20 (3.7)	204 (4.1)	21 (3.7)	27 (4.8)
Parity:				
Multiparous	338 (62.4)	2789 (55.4)	115 (19.8)	174 (30.6)
Primiparous	204 (37.6)	2245 (44.6)	465 (80.5)	394 (69.4)
Gestational age at birth				
<37 weeks	57 (10.5)	196 (3.9)	22 (3.8)	86 (15.1)
≥37 weeks	485 (89.5)	4838 (96.1)	558 (96.2)	482 (84.9)
Induced	13 (2.4)	813 (16.2)	143 (24.7)	188 (33.1)
Epidural Analgesia	67 (12.4)	1322 (26.3)	342 (59.0)	283 (49.8)
Birth weight:				
<2500 g	45 (8.3)	114 (2.3)	12 (2.1)	72 (12.7)
2500–4000 g	478 (88.4)	4746 (94.6)	552 (95.8)	459 (80.8)
≥4000 g	18 (3.3)	157 (3.1)	12 (2.1)	37 (6.5)
Apgar score:				
<7 after 5 min	4 (0.7)	34 (0.7)	20 (3.5)	16 (2.8)
≥7 after 5 min.	537 (99.3)	4948 (99.3)	556 (96.5)	547 (97.2)
Twins	43 (8.1)	74 (1.5)	17 (3.0)	22 (3.9)

doi:10.1371/journal.pone.0087579.t003

delivery; hence, the observed associations may be seen as results of the study as an intervention. However, if this were the case, we would have likely seen increases in the estimated associations in all types of abuse.

Another important strength of our study is that we used a previously validated instrument to measure abuse. The validity of the NorAQ compared to clinical interviews and other instruments has been shown to be high [20]. Still women may have opted not to report abuse. This may have biased the estimated association towards zero. As this is a longitudinal study, a recall bias based on outcome is unlikely. Our study did not include questions on the onset, length of time and frequency of the abuse, nor who the perpetrator was. The influence of these factors could therefore not be studied. Our findings are based upon women from selected European countries, and may not be generalized to other locations with very different health care system.

Some of our findings are in conflict with other studies. A cohort study from Trondheim found that childhood abuse, whether it was physical or sexual, was associated with an increased risk of interventions during childbirth, both CS and instrumental vaginal delivery [13]. This study made no distinction between primiparous and multiparous women [13]. However, the lack of associations between childhood abuse and operative delivery agrees with the

findings of a large population based study from Norway, which found no association between childhood abuse and CS before labor and only a slight increase for CS during labour [14]. The lack of association between childhood abuse and mode of delivery in our study could be due to low power in our study due to few cases. Alternatively, it may be the result of effective psychosocial counselling. An increased recognition and optimal intervention for depression and fear of delivery may prevent interventions. A Norwegian study including women attending a specialised clinic aimed at caring for women with fear of childbirth, reported that sexual abuse in the form of rape after the age of 16 years was associated with a major increase in risk for CS [15]. This is in agreement with our results. However, theirs was a selected population with whom mode of delivery was actively discussed as part of the consultation and hence the association may have been the consequence of an active choice during the consultation.

Sexual abuse is likely to affect pregnant women more than other types of abuse since the female reproductive organs are involved both in sexual acts and in giving birth [12,28]. For this reason, the woman and/or the obstetrical staff may be anxious about a vaginal delivery, perhaps more so than for a history of physical or emotional abuse. An obstetrician who is aware of such a history may be more inclined to grant a CS on maternal request without

Table 4. The association between (Adjusted OR*) different types of abuse and operative delivery for multiparous women (n = 3416) in the Bidens study.

	Abuse	Elective CS	Elective CS	Operative vaginal	Operative vaginal	Emergency CS	Emergency CS
	n (%)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)
No abuse	2214 (64.8)	9.8	1	3.6	1	4.9	1
Any abuse	1202 (35.2)	10.1	1.05 (0.83–1.33)	3.0	0.83 (0.56–1.23)	5.4	1.11 (0.80–1.53)
Any abuse <18	797 (23.3)	10.3	1.08 (0.82–1.41)	3.1	0.88 (0.55–1.39)	5.5	1.13 (0.78–1.64)
Any abuse ≥18	704 (20.6)	9.7	0.98 (0.73–1.31)	2.8	0.79 (0.48–1.31)	5.0	1.00 (0.67–1.49)
Emotional abuse	638 (18.7)	9.7	1.00 (0.74–1.35)	2.2	0.61 (0.34–1.09)	6.3	1.30 (0.89–1.91)
Emotional abuse <18	385 (11.3)	10.6	1.12 (0.78–1.61)	2.3	0.66 (0.33–1.33)	6.2	1.31 (0.82–2.10)
Emotional abuse ≥18	369 (10.8)	8.7	0.86 (0.58–1.28)	2.4	0.67 (0.33–1.36)	5.7	1.15 (0.70–1.87)
Physical abuse	653 (19.1)	11.3	1.24 (0.93–1.65)	3.5	1.03 (0.64–1.66)	6.9	1.51 (1.05–2.19)
Physical abuse <18	328 (9.6)	11.6	1.28 (0.88–1.86)	3.0	0.89 (0.46–1.75)	7.3	1.57 (0.97–2.52)
Physical abuse ≥18	383 (11.2)	10.7	1.13 (0.79–1.62)	3.4	0.99 (0.54–1.80)	6.0	1.30 (0.81–2.08)
Sexual abuse	561 (16.4)	10.5	1.09 (0.80–1.49)	2.7	0.74 (0.42–1.30)	4.8	0.97 (0.62–1.51)
Sexual abuse <18	398 (11.6)	10.1	1.05 (0.73–1.50)	2.8	0.77 (0.40–1.46)	4.8	0.96 (0.58–1.60)
Sexual abuse ≥18	221 (6.5)	11.3	1.14 (0.73–1.78)	2.7	0.77 (0.33–1.80)	4.5	0.91 (0.46–1.79)

*Adjusted for age, twin pregnancy, gestational age less than 37 weeks, and country of residence. Compared to women not reporting any abuse.
doi:10.1371/journal.pone.0087579.t004

any other medical indication, as our observed association indicate. In a situation where there is uncertainty about mode of delivery, knowledge of a woman's history of abuse and/or her wish for CS may influence the decision towards a CS with a medical indication. In our study, we had no knowledge about whether women revealed their history and/or preference for CS at any stage before delivery. There is no consensus about the optimal mode of delivery for women with a traumatic history of adult sexual abuse, nor do we know how alternative approaches may

affect the mother-infant relationship and the long-term psychological effects of delivery/childbirth. However, our results suggest that efforts should be made to reduce suffering from the experienced abuse as current suffering increased the risk of not only elective but also emergency CS.

Women with a previous delivery and a history of abuse did not have an increased risk of operative delivery associated with the abuse. A possible explanation could be that victims of adult sexual abuse who either had an operative delivery and/or suffered a poor

Table 5. The association (Adjusted OR*) between different types of abuse and mode of delivery for primiparous women (n = 3308) in the Bidens study.

	Abuse	Elective CS	Elective CS	Operative vaginal	Operative vaginal	Emergency CS	Emergency CS
	n (%)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)
No abuse	2187 (66.1)	5.9	1	13.8	1	11.5	1
Any abuse	1121 (33.9)	6.7	1.22 (0.90–1.66)	14.5	1.11 (0.97–1.37)	12.7	1.16 (0.92–1.45)
Any abuse <18	770 (23.3)	5.6	1.04 (0.72–1.51)	14.7	1.13 (0.89–1.44)	13.0	1.21 (0.94–1.56)
Any abuse ≥18	605 (18.3)	8.6	1.45 (1.02–2.06)	13.2	0.97 (0.74–1.27)	12.6	1.09 (0.82–1.45)
Emotional abuse	623 (18.8)	7.4	1.38 (0.96–1.98)	15.1	1.21 (0.93–1.57)	13.3	1.24 (0.94–1.64)
Emotional abuse <18	422 (12.7)	6.6	1.28 (0.82–1.98)	14.2	1.15 (0.84–1.56)	14.2	1.37 (1.00–1.88)
Emotional abuse ≥18	316 (9.5)	8.5	1.50 (0.96–2.36)	13.9	1.07 (0.76–1.53)	13.3	1.17 (0.82–1.69)
Physical abuse	566 (17.1)	6.7	1.21 (0.82–1.78)	14.1	1.03 (0.78–1.36)	10.2	0.93 (0.68–1.27)
Physical abuse <18	339 (10.2)	5.6	1.05 (0.63–1.76)	14.2	1.07 (0.76–1.50)	10.6	1.00 (0.69–1.47)
Physical abuse ≥18	297 (9.0)	8.8	1.45 (0.92–2.30)	13.8	0.97 (0.68–1.40)	9.4	0.81 (0.53–1.23)
Sexual abuse	495 (15.0)	7.7	1.42 (0.96–2.10)	14.1	1.07 (0.80–1.43)	12.7	1.19 (0.88–1.61)
Sexual abuse <18	342 (10.3)	5.5	1.00 (0.60–1.67)	12.9	0.93 (0.66–1.32)	12.0	1.07 (0.75–1.54)
Sexual abuse ≥18	200 (6.0)	11.0	2.12 (1.28–3.49)	16.0	1.31 (0.87–1.98)	14.0	1.39 (0.90–2.16)

*Adjusted for age, twin pregnancy, gestational age less than 37 weeks, and country of residence. Compared to women not reporting any abuse.
doi:10.1371/journal.pone.0087579.t005

Table 6. The association between adult sexual abuse and mode of delivery for primiparous women (n = 3308) in the Bidens study.

Exposure	n	Elective CS		Operative vaginal delivery		Emergency CS	
		%	Adjusted OR (95% CI)	%	Adjusted OR (95% CI)	%	Adjusted OR (95% CI)
No abuse	2187	5.9	1	13.8	1	11.5	1
Any adult sexual abuse*	200	11.0	2.12 (1.28–3.49)	16.0	1.31 (0.87–1.98)	14.0	1.39 (0.90–2.16)
Adult sexual abuse*							
mild	46	15.2	2.21 (0.93–5.26)	6.5	0.42 (0.13–1.38)	8.7	0.69 (0.24–1.99)
moderate	41	9.8	1.77 (0.57–5.52)	19.5	1.72 (0.75–3.92)	17.1	1.83 (0.76–4.40)
severe	113	9.7	2.15 (1.10–4.23)	18.6	1.66 (1.00–2.77)	15.0	1.60 (0.91–2.79)
Adult sexual abuse and:							
adult physical*	70	12.9	2.88 (1.34–6.19)	17.1	1.52 (0.78–2.95)	14.3	1.61 (0.79–3.30)
adult emotional*	59	13.6	2.93 (1.30–6.60)	20.3	1.84 (0.93–3.63)	11.9	1.27 (0.55–2.93)
adult emotional and physical*	38	10.5	2.69 (0.88–8.21)	23.7	2.34 (1.04–5.26)	15.8	2.02 (0.79–5.16)
current suffering*	37	13.5	4.07 (1.46–11.3)	16.2	1.86 (0.72–4.74)	21.6	2.97 (1.26–6.98)
child sexual abuse*	47	6.4	1.09 (0.32–3.69)	12.8	0.93 (0.38–2.27)	12.8	1.08 (0.44–2.65)
recent experience†	12	16.7	3.52 (0.73–16.8)	0	0	8.3	0.82 (0.10–6.48)

*Adjusted age, twin pregnancy, gestational age less than 37 weeks, and country of residence.

†Due to few cases adjusted only for age.

Compared to women not reporting any abuse.

doi:10.1371/journal.pone.0087579.t006

birth experience chose not to become pregnant again and hence they were not among the multiparous women in our study. It is also possible that the previous birth experience was a healing experience as described by Simpkin and Klaus [29].

An obvious possible reason for an increase in emergency CS among women reporting physical abuse would be placental abruption after trauma. This indication was not among the options in the outcome sheet in our study. The most common CS indications for these women were fetal distress and other medical indications, which could be related to direct physical trauma to the abdomen. However, when we added previous CS into the analysis, the association was clearly attenuated, indicating that this factor played a major role. The CS in the previous pregnancy could have been due to abuse prior to that. Our results indicated that a history of abuse has limited impact on mode of delivery for women with a previous birth experience. As in obstetrics in general, the method of the first birth has a great influence on subsequent mode of delivery [30].

Conclusions

Among primiparous women sexual abuse experienced as an adult was associated with increased elective caesarean sections. Primiparous women with a history of adult sexual abuse were also

more likely to have a non-obstetrically indicated CS. Identifying a woman's sexual abuse status during pregnancy may influence decision-making regarding mode of delivery.

Acknowledgments

We are greatly indebted to all women participating in the study. Likewise to all the staff actively participating and facilitating the study in all the different study sites. We would like to thank Professor Svein Ove Samuelsen for statistical advice.

Author Contributions

Conceived and designed the experiments: BS JC MT. Performed the experiments: AMS H. Kristjansdottir AMW M. Lukasse M. Laanpere ASVP H. Karro AT MT ELR BS TS. Analyzed the data: Mirjam L. Contributed reagents/materials/analysis tools: AMS H. Kristjansdottir AMW M. Lukasse M. Laanpere ASVP H. Karro AT MT ELR BS TS JC. Wrote the paper: BS M. Lukasse ELR TS H. Karro. Commented on drafts and approved the final manuscript: AMS H. Kristjansdottir AMW M. Lukasse M. Laanpere ASVP H. Karro AT MT ELR BS TS JC. Developed the questionnaire: BS M. Lukasse ELR JC H. Karro H. Kristjansdottir M. Laanpere AMS AT MT ASVP AMW TS. Finalized the research protocol: BS M. Lukasse ELR JC H. Karro H. Kristjansdottir M. Laanpere AMS AT MT ASVP AMW TS. Collected data: AMS AMW M. Lukasse M. Laanpere ASVP.

References

1. Oppenheimer L (2007) Diagnosis and management of placenta previa. *J Obstet Gynaecol Can* 29: 261–273.
2. Souza JP, Gulmezoglu A, Lumbiganon P, Laopaiboon M, Carroli G, et al. (2010) Caesarean section without medical indications is associated with an increased risk of adverse short-term maternal outcomes: the 2004–2008 WHO Global Survey on Maternal and Perinatal Health. *BMC Med* 8: 71.
3. Lavender T, Hofmeyr GJ, Neilson JP, Kingdon C, Gyte GM (2012) Caesarean section for non-medical reasons at term. *Cochrane Database Syst Rev* 3: CD004660.
4. Chambliss LR (2008) Intimate partner violence and its implication for pregnancy. *Clin Obstet Gynecol* 51: 385–397.
5. Fanslow J, Silva M, Robinson E, Whitehead A (2008) Violence during pregnancy: associations with pregnancy intendedness, pregnancy-related care, and alcohol and tobacco use among a representative sample of New Zealand women. *Aust N Z J Obstet Gynaecol* 48: 398–404.
6. Jun HJ, Rich-Edwards JW, Boynton-Jarrett R, Wright RJ (2008) Intimate partner violence and cigarette smoking: association between smoking risk and psychological abuse with and without co-occurrence of physical and sexual abuse. *Am J Public Health* 98: 527–535.
7. Oxford CM, Ludmir J (2009) Trauma in pregnancy. *Clin Obstet Gynecol* 52: 611–629.
8. Brown HL (2009) Trauma in pregnancy. *Obstet Gynecol* 114: 147–160.
9. Keeshin BR, Cronholm PF, Strawn JR (2012) Physiologic changes associated with violence and abuse exposure: an examination of related medical conditions. *Trauma Violence Abuse* 13: 41–56.

10. Lukasse M, Vangen S, Oian P, Schei B (2011) Fear of childbirth, women's preference for cesarean section and childhood abuse: a longitudinal study. *Acta Obstet Gynecol Scand* 90: 33–40.
11. Lukasse M, Vangen S, Oian P, Kumle M, Ryding EL, et al. (2010) Childhood abuse and fear of childbirth—a population-based study. *Birth* 37: 267–274.
12. Rhodes N, Hutchinson S (1994) Labor experiences of childhood sexual abuse survivors. *Birth* 21: 213–220.
13. Heimstad R, Dahloe R, Laache I, Skogvoll E, Schei B (2006) Fear of childbirth and history of abuse: implications for pregnancy and delivery. *Acta Obstet Gynecol Scand* 85: 435–440.
14. Lukasse M, Vangen S, Oian P, Schei B (2010) Childhood abuse and caesarean section among primiparous women in the Norwegian Mother and Child Cohort Study. *BJOG* 117: 1153–1157.
15. Nerum H, Halvorsen L, Oian P, Sorlie T, Straume B, et al. (2010) Birth outcomes in primiparous women who were raped as adults: a matched controlled study. *BJOG* 117: 288–294.
16. Stenson K, Heimer G, Lundh C, Nordstrom ML, Saarinen H, et al. (2003) Lifetime prevalence of sexual abuse in a Swedish pregnant population. *Acta Obstet Gynecol Scand* 82: 529–536.
17. van der Hulst LA, Bonsel GJ, Eskes M, Birnie E, van Teijlingen E, et al. (2006) Bad experience, good birthing: Dutch low-risk pregnant women with a history of sexual abuse. *J Psychosom Obstet Gynaecol* 27: 59–66.
18. Nerum H, Halvorsen L, Straume B, Sorlie T, Oian P (2012) Different labour outcomes in primiparous women that have been subjected to childhood sexual abuse or rape in adulthood: a case-control study in a clinical cohort. *BJOG*.
19. Wijma B, Schei B, Swahnberg K, Hilden M, Offerdal K, et al. (2003) Emotional, physical, and sexual abuse in patients visiting gynaecology clinics: a Nordic cross-sectional study. *Lancet* 361: 2107–2113.
20. Swahnberg IM, Wijma B (2003) The NorVold Abuse Questionnaire (NorAQ): validation of new measures of emotional, physical, and sexual abuse, and abuse in the health care system among women. *Eur J Public Health* 13: 361–366.
21. Swahnberg K, Schei B, Hilden M, Halmesmaki E, Sidenius K, et al. (2007) Patients' experiences of abuse in health care: a Nordic study on prevalence and associated factors in gynecological patients. *Acta Obstet Gynecol Scand* 86: 349–356.
22. Wijma K, Wijma B, Zar M (1998) Psychometric aspects of the W-DEQ: a new questionnaire for the measurement of fear of childbirth. *J Psychosom Obstet Gynaecol* 19: 84–97.
23. Eberhard-Gran M, Eskild A, Samuelsen SO, Tambs K (2007) A short matrix-version of the Edinburgh Depression Scale. *Acta Psychiatr Scand* 116: 195–200.
24. Ellsberg M, Heise L (2002) Bearing witness: ethics in domestic violence research. *Lancet* 359: 1599–1604.
25. Lukasse M, Schei B, Vangen S, Oian P (2009) Childhood abuse and common complaints in pregnancy. *Birth* 36: 190–199.
26. Silverman JG, Decker MR, Reed E, Raj A (2006) Intimate partner violence victimization prior to and during pregnancy among women residing in 26 U.S. states: associations with maternal and neonatal health. *Am J Obstet Gynecol* 195: 140–148.
27. Garcia-Moreno C, Jansen HAFM, Elsberg M, Heise L, Watts C (2005) WHO Multi-country Study on Women's Health and Domestic Violence against Women Initial results on prevalence, health outcomes and women's responses WHO, editor. Geneva, Switzerland.
28. Hobbins D (2004) Survivors of childhood sexual abuse: implications for perinatal nursing care. *J Obstet Gynecol Neonatal Nurs* 33: 485–497.
29. Simkin P, Klaus PH (2004) When Survivors Give Birth: Understanding and Healing the Effects of Early Sexual Abuse on the Childbearing Woman. Seattle, Washington: Classis Day. 450 p.
30. Cunningham FG, Bangdiwala SI, Brown SS, Dean TM, Frederiksen M, et al. (2010) NIH consensus development conference draft statement on vaginal birth after cesarean: new insights. *NIH Consens State Sci Statements* 27: 1–42.